

# LA-UR-12-22318

Approved for public release; distribution is unlimited.

Title:	MCNP6 Cosmic-Source Option
Author(s):	McKinney, Gregg W Armstrong, Hirotatsu James, Michael R Clem, John Goldhagen, Paul
Intended for:	ANS Annual Meeting, 2012-06-24/2012-06-28 (Chicago, Illinois, United States)



#### Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

## **ABSTRACT**

MCNP is a Monte Carlo radiation transport code that has been under development for over half a century. Over the last decade, the development team of a high-energy offshoot of MCNP, called MCNPX, has implemented several physics and algorithm improvements important for modeling galactic cosmic-ray (GCR) interactions with matter. In this presentation, we discuss the latest of these improvements, a new Cosmic-Source option, that has been implemented in MCNP6.

# MCNP6 Cosmic-Source Option

---

Gregg McKinney<sup>1</sup>, James Armstrong<sup>1</sup>, Mike James<sup>1</sup>,  
John Clem<sup>2</sup>, Paul Goldhagen<sup>3</sup>

(1) LANL, International & Nuclear System Engineering Group (D-5)

(2) University of Delaware, Bartol Research Institute

(3) DHS, National Urban Security Technology Laboratory

*ANS Annual Meeting, Chicago, IL, June 24-28, 2012*


# Outline

---


- **Introduction**
- **Cosmic-source option**
- **Benchmark results**
- **Terrestrial backgrounds**
- **Conclusions**

US Department of Homeland Security, Domestic Nuclear Detection Office, under competitively awarded contract/ IAA HSHQDC-09-X-00190. This support does not constitute an express or implied endorsement on the part of the Government.


# Neutron & gamma GCR background spectra show a strong dependence on longitude, latitude, and altitude



# GCR composition – initial MCNP6 implementation includes protons and alphas




# GCR induced neutron and photon spectra at sea level (Livermore, CA)



# Collaborations are underway to provide a predictive world-wide background capability

## Cosmic Radiation



“Sky Maps” of protons, alphas, etc. versus long. & lat.  
(U of Del. – John Clem)

Detailed cosmic ray model  
(300 layers of atmosphere)  
(MCNPX)

Cosmic neutron benchmark  
measurements from Long  
Dwell program  
(NUSTL – Paul Goldhagen)

NORM Spectra - K, Rb, Th, U  
(MCNPX)

Terrestrial Radiation



# Cosmic-source option – two GCR spectra (interplanetary, geomagnetic) and 3 new SDEF source keywords

---

- **Built-in spectra**
  - Historic (PRL/Lal, 1980)
  - Modern (UoD/Clem, 2006)
- **User interface**
  - PAR keyword enhanced
  - New keyword LOC
  - New keyword DAT
- **Benchmarking (NUSTL/Goldhagen collaboration)**
  - Atmosphere (primary spectra, soon)
  - Atmosphere (secondary spectra ~120°W, June 1997)
  - Ground (NY, Nov. 2001, soon; CA, Nov. 2006)

# Interplanetary spectrum proposed by Devendra Lal, combined with rigidity cutoffs

---

- **Intended for interplanetary use**

- See Apollo 17 benchmark paper (McKinney et al., JGR, 2006)

- **Analytic form of the  $4\pi$  differential spectra**

$$g(T, \phi) = A T (T + 2E_0) (T + m + \phi)^{-\gamma} / [(T + \phi) (T + 2E_0 + \phi)]$$

$g$  – particles/cm<sup>2</sup>-s-MeV

$T$  = kinetic energy per nucleon (MeV/n)

$\phi$  = solar modulation potential (MV)

$E_0$  = rest energy of a nucleon (MeV)

$m = a \exp(-bT)$


$A, a, b, \gamma$  are fitting parameters (different for each particle)

- **Apply Clem rigidity cutoffs for geomagnetic use**

- Truncates low-energy portion of Lal spectra

# Geomagnetic spectra proposed by John Clem, uses skymaps describing rigidity parameters

- **Intended for geomagnetic use**
  - Has a built-in interplanetary spectrum (zero rigidity)
- **Skymaps generated on a geomagnetic grid**
  - Provides rigidity cutoffs versus polar & azimuthal angle
  - Separate set of maps for various altitudes
  - Monte Carlo code provides light & heavy ion spectra ( $Z > 2$ )
- **MCNP6 includes a link to the BRI UoD code**
  - Skymap data condensed on  $5^\circ$  lat.,  $20^\circ$  long. grid (at 65 km)



# MCNP6 user interface – three new SDEF keywords

**Clem Cosmic Source for Solar Max.**

```
1 0 -1 2 imp:n=1
2 0 -1 -2 imp:n=1
3 0 1 imp:n=0
```

```
1 rpp -1000 1000 -1000 1000 -1000 1000
2 pz 999
```


```
mode n h a
SDEF X=d1 Y=d2 Z=999.1 VEC=0 0 -1
      PAR=CR DAT=1 1 1987 LOC=80 -120 65
si1 -1000 1000
sp1 0 1
si2 -1000 1000
sp2 0 1
phys:h,a 1e6
nps 1000000
print
f11:h 2
f12:h 2
fm12 4E6
f21:a 2
f22:a 2
fm22 4E6
e0 1 349ilog 1e6
c dbcn 31j 1 $ Use Lal spectra
```

**Description of SDEF keywords.**

Keyword	Values	Description
PAR	[-]cr	All cosmic particles
	[-]ch	Cosmic protons
	[-]ca	Cosmic alphas
DAT	M	Month (1-12)
	D	Day (1-31)
	Y	Year (4 digit)
LOC	P	Latitude (-90 to 90; S to N)
	A	Longitude (-180 to 180; W to E)
	H	Altitude (km)


# Top-of-the-atmosphere proton and alpha spectra at various longitudes and latitudes – solar minimum

## Solar Minimum (1987)




# Top-of-the-atmosphere proton and alpha spectra at various longitudes and latitudes – solar maximum


## Solar Maximum (1981)



# Neutron spectra compared to measured data – NASA flights over western Canada, CA, and west of Mexico



# Neutron spectra compared to measured data – Livermore, CA, November 2006








# Neutron spectra compared to measured data – ongoing New York, NY, November 2001 and 2012



# Terrestrial radionuclide concentrations will be taken from USGS maps – initially using a 10°x10° long./lat. grid



# Simple example of radionuclides in a wet porous soil (Livermore, CA) – tally photon flux above the ground

## Terrestrial background for 37N 120W

```
1 1 -1.4000 1 -2 -5 imp:p=1
2 2 -0.00120698 2 -3 -5 imp:p=1
3 2 -0.00120698 3 -4 -5 imp:p=1
4 0 5:-1:4 imp:p=0
```

```
1 pz 0.0
2 pz 100.0
3 pz 200.0
4 pz 10000.0
5 cz 1000.0
```

mode p

act fission=none nonfiss=p dg=lines

m1 8016 -.58457 \$ Wet porous soil

```
14028 -.18944
14029 -.00962
14030 -.00634
13027 -.06026
26054 -.00217
26056 -.03400
26057 -.00079
26058 -.00010
1001 -.03013
20000 -.02691
11023 -.02098
19000 -.01920
12000 -.01549
```

```
19040 -2e-6 $ 2 ppm
90232 -5e-6 $ 5 ppm
92238 -2e-6 $ 2 ppm
m2 $ Air with 60% humidity
7014 -.729981
8016 -.256290
18000 -.012374
1001 -.001360
sdef par=sp wgt=2.5e8
pos=0 0 0 axs=0 0 1 rad=d1 ext=d2
si1 0 1000
sp1 -21 1
si2 0 100
sp2 0 1
f4:p 2
e4 1e-3 299log 10
nps 1000000
print
```




# Terrestrial photon flux combined with cosmic photon flux on a 10°x10° grid within the MCNP6 “background.dat” file

Neutron flux from Goldhagen JEDEC No. 89A (2006)		Photon flux from MCNPX	
90.0	0.0	0.0	183
flux (n/cm2/s)	0.1193E-01	flux (p/cm2/s)	2.0000E+00
1.0593E-10	2.1270E+02	1.0354E-03	2.5895E-05
1.1885E-10	2.3849E+02	1.1087E-03	1.3244E-05
1.3335E-10	2.6743E+02	1.1872E-03	5.5672E-05
1.4962E-10	3.0003E+02	1.2712E-03	2.5714E-05
1.6788E-10	3.3624E+02	1.3612E-03	1.1886E-05
1.8836E-10	3.7715E+02	1.4576E-03	5.1054E-06
2.1135E-10	4.2306E+02	1.5608E-03	3.8590E-05
...			
		1.6937E+01	7.4518E-05
		1.7575E+01	5.2691E-05
		1.8237E+01	6.4417E-05
		1.8924E+01	5.0422E-05
		1.9637E+01	4.4057E-05
		2.0797E+01	5.4823E-05
Neutron flux from Goldhagen JEDEC No. 89A (2006)		Photon flux from MCNPX	
0.0	0.0	0.0	0
flux (n/cm2/s)	0.6995E-02	flux (p/cm2/s)	2.0000E+00
Neutron flux from Goldhagen JEDEC No. 89A (2006)		Photon flux from MCNPX	
0.0	10.0	0.0	0
flux (n/cm2/s)	0.6914E-02	flux (p/cm2/s)	2.0000E+00
Neutron flux from Goldhagen JEDEC No. 89A (2006)		Photon flux from MCNPX	
0.0	20.0	0.0	0
flux (n/cm2/s)	0.6828E-02	flux (p/cm2/s)	2.0000E+00
...			

Terrestrial  
+  
Cosmic

# Predicted background spectra for all particles at sea level (Livermore, CA)

SNLL (37°N, 120°W, ~0 km), Nov. 2006 (solar min.)



# Conclusions

---

- **MCNP6 can provide realistic world-wide cosmic & terrestrial background spectra, which can be combined with other standard sources**
- **Benchmarking of these capabilities will continue**
- **Incorporation of cosmic & terrestrial spectra into the “background.dat” file will continue (10°x10° grid) to facilitate its use in analyses of nuclear systems**
- **Future upgrades include:**
  - Heavy-ion GCR
  - Altitude dependence
  - Improved interpolation of “background.dat” spectra
  - Additional cosmic spectra (proton, alpha, muon, pion, etc.)
  - Improved geomagnetic grid (more data where changes are large)